701254

Hay 27. 1947

P. O. Box 1663 Sante Fe, Hew Mexico

Door Wrights

You will receive another shipment of specimens from Hp9 within a few days. These include five (5) 24 hoursuring specimens and two bottles containing the feces collected between carmine markers given at 8:00 A.M., 5/15/47 and 8:00 A.M., 5/21/47. These specimens have been collected during the intravenous administration of sodium citrate solution. The citrate was given intravenously in a concentration of 1.5% dissolved in physiological caline. The dates of injection, interval over which injected and total amount are given in the following tobulations:

Date	Time Taken for Injection	Amount gas.
5/16/47	120 nin-	10.0
5/17/47	145	15.0
5/15/47	160	15-0
5/1 <b>9/47</b>	180	15.0
5/20/47	160	15.0

We were rather cautious about the amount given since some data in the medical literature suggested that large amounts of sodium citrate are toxic. In paper by Bruneau and Graham, Arch. Surg., 1985, 47, 319, it is toxic stated that 15 gm. of sodium citrate may be toxic if given rapidly. We did not notice any adverse effect giving it slowly as indicated. I presume considerably larger amounts would be tolorated but perhaps not by hip who is a difficult suggest in many respects.

Sincerely yours,

Sequel E. Sassett

SEE: Phb

Classification Cancelled

Or Changest To

By Authority Of Diver Castilian 13

By K. Ho teripleb Date 5/4

0004125